

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/557	A1	(11) International Publication Number: WO 98/50039 (43) International Publication Date: 12 November 1998 (12.11.98)
(21) International Application Number: PCT/GR98/00012 (22) International Filing Date: 22 April 1998 (22.04.98) (30) Priority Data: 970100172 6 May 1997 (06.05.97) GR (71)(72) Applicants and Inventors: KANAKARIS, Panagiotis [GR/GR]; 33 Koletti Street, GR-106 77 Athens (GR). KAROUZAKIS, Petros [GR/GR]; 9 Bakou Street, GR-115 24 Athens (GR).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: USE OF MISOPROSTOL OR/AND MISOPROSTOL ACID FOR PREPARING DRUG IN ORDER TO CURE ERECTILE DYSFUNCTION (57) Abstract The invention relates to using an already known pharmaceutical substance, misoprostol or/and its first metabolite (misoprostol acid) whose structural formulas are presented in pages 3 and 5 (Figures I and II), for preparing a drug of external use destined: a) to cure erectile dysfunction and b) as accessory means in the diagnosis of erectile dysfunction. Misoprostol or/and misoprostol acid according to the described method are applied externally on the glans of the penis, the prepuce and the urethra, are absorbed and cause topical vasodilation resulting in a hard and durable erection for patients suffering from disorders of erection, due to vascular or other causes.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

USE OF MISOPROSTOL OR/AND MISOPROSTOL ACID FOR PREPARING DRUG
• IN ORDER TO CURE ERECTILE DYSFUNCTION

The invention relates to the use of an already known
5 pharmaceutical substance, misoprostol as well as its first
metabolite, misoprostol acid, for the preparation of a drug for
external use which is destined to cure erectile dysfunction.
Nowadays the pharmaceutical treatment of erectile dysfunction
-except of the cases of hormonal insufficiency, which are
10 generally rare and in which a suitable substitution therapy
is followed-includes mainly the use of intracavernosous
injections consisting in the direct injection of vasodilatory
drugs (papaverine, phentolamine and alprostadil) into the
corpora cavernosa of the penis (Campell's Urology, ed. W. B.
15 Saunders Company, 6th edition, volume III, p. 3055-3057).
Although this method is the most efficient and perhaps the
only scientifically acceptable, it has the serious
disadvantage of the form (injection) as well as the manner of
administration intracavernosal.
20 Yohimbin, an α_2 -adrenergic inhibitor, is administered per os,
however the efficiency of this old method is doubtful
(Campell's Urology, ed. W. B. Saunders Company, 6th edition,
volume III, p. 3053).
In former times the topical application of a nitroglycerin
25 paste had been proposed (Claes et al 1989), but the method
was not therapeutically applied because of doubtful
efficiency and serious side-effects (Campell's Urology,
ed. W. B. Saunders Company, 6th edition, volume III, p. 3053).
The topical application of prostaglandin E₁ (or alprostadil)

in the form of an endourethral gel or stick as a means of
• limited efficiency in the therapy of male impotence of a
vascular cause (International Journal of Impotence Research,
Stocton ed., vol. 7, September 1995, supplement I, p. 05-06) was
5 recently proposed, we must note that the discovery of
vasodilatory drugs with sufficient transcutaneous absorption
or the use of methods (e.g. ionophoresis) which can reinforce
the penetration of such drugs through the skin of the mucosal
membranes, inside the corpora cavernosa of the penis has for
10 long attracted the interest of many research workers
(Campell's Urology, ed. W. B. Saunders Company, 6th ed., vol. III,
p. 3057)

Up to day a common denominator of the methods destined for
external application is mainly the low efficiency combined
15 with increased therapy cost and the apparition of more or
less serious side-effects.


The present method aims at the removal of the drawbacks of
the above methods using misoprostol in the symptom therapy of
male impotency. Misoprostol is the general name of a synthetic
20 prostaglandin belonging to the E₁ series (PGE₁ analogs).

Synthesis: P. W. Collins, R. Pappo, Belgian patent 827.127,
American patent 3.965.143 (The Merck Index, ed. Merck & Co.
Inc, 11th edition, 1989, p. 6128).

Its chemical name is (11a,13E)-(±)-11,16-Dihydroxy-16-methyl-
25 9-oxoprost-13-en-1-oic acid methyl ester or (±)-(methyl)-
(1R, 2R, 3R)-3-hydroxy-2-[(E)-(4RS)-4-hydroxy-4-methyl-1-
octenyl]-5-oxocyclopentaneheptanoate or (±)-15-deoxy-(16RS)-
16-hydroxy-16-methyl-PGE₁ methyl ester.

It is consisted of 4 stereoisomers in about equal proportions
 • [(+)&(-) enantiomers of 16R- and 16S-forms]. (The Merck Index,
 11th edition, 1989, p. 6128). The empirical formula is C₂₂H₃₈O₅.
 Its structural formula appears in Fig. I.

5



15

Figure I.

Compared with other prostaglandins of group E₁ and especially
 alprostadil, misoprostol bears a methyl group (-CH₃) on the
 20 carbon atom of position 16.

According to a method which relates the biological action of
 various medicament molecules to its chemical structure
 (Method of Minimum Stereochemical Difference, "Planning of
 drugs", P. Kourounakis-E. Rekka, ed. Graphical
 25 Arts, Thessaloniki, 1992, p. 152) it appears that due to this
 group we have a big penetration of misoprostol in the
 underlying tissues and a local vasodilation which cures
 erection dysfunctions. Misoprostol is used today orally as

antiulcer drug (Physicians Desc Reference, PDR, ed. Medical Economics Data, Production Company at Montrale 48th edition, 1994, P. 2197-2199).

- In particular it is administered for the prevention of gastric ulcer to patients who take non-steroid antiinflammatory drugs. It is available in the countries of Europe and U. S. A. by Searle Company under the commercial name Cytotec®. In none country is the drug mentioned as suitable for male impotence nor are there any relevant reports on the international bibliography. On a contrary amongst the undesirable effects in oral therapy with misoprostol is male impotence (Physicians Desc Reference, ed. Medical Economics Data, Production Company at Montrale, 48th edition, 1994, p. 2197-2199).
- 15 In spite of the fact that misoprostol exerts reduced vasodilatory action compared with alprostadil when administered intracavernously, it creates a stronger local vasodilation and consequently greater increase in blood flux when applied externally to the penis and the urethral
- 20 mucosa. Due to the topical vasodilation we have a sufficiently long and durable erection. On the other hand due to the fact that the degree of response to a small dose of misoprostol (durability and hardness of erection) depends on the physiological condition and function of the penile
- 25 vessels, misoprostol can be used as accessory diagnostic means (instead of papaverine or alprostadil) in the "Doppler" method or the cavernosometry, for the determination of the extent and kind of vascular damage (about the use of

vasodilatory drugs as accessory diagnostic means in the

• "Doppler" method or cavernosometry, see Erektile

Impotenz, ed. Enke, p. 68-77 & p. 88-110).

Equally strong topical vasodilatory action after external

5 application is exerted by the hydrolysis product of

misoprostol, (misoprostol acid) which anyway constitutes the

first misoprostol metabolite after its introduction in the

organism (see Fig. II).

10

15

20




Figure II.

Last because of the intense topical vasodilatory action of

misoprostol and the corresponding free acid, the two

pharmaceutical molecules reinforce the absorption of other

25 vasoactive substances (e.g. alprostadil) resulting in the

occurrence of synergic action.

Misoprostol can be dissolved in water and its compatibility

with excipients provides the opportunity of production of a

variety of simple pharmacotechnical forms for external use, which are at the same time very well tolerated by the skin and the mucousa. From the above mentioned description it appears that the most

- 5 serious advantage of the method is the manner of administration of the drug (external in combination with the lack of undesirable action in the suggested doses or/and the proposed pharmacotechnical forms) the relatively low cost and especially the most satisfactory result together with
- 10 corresponding methods.

Amongst the probable methods of application, most advantageous is a synthesis in the gel form of relatively low viscosity which contains 0,9 % w/v misoprostol in the methylform of methylester and/or free acid, a complexforming means, as

- 15 1,6% w/v α -cyclodextrine and substances suitable for the formation of a gel e.g. hydroxypropyl methylcellulose "3000" 2% w/v, propylene glycol 10% v/v and Water to 100 ml. The gel contains 9 mg of active substance per ml.

- Method of application: 0,1-0,25 ml (or more, depending on
- 20 response) are pasted or spread on the glans of penis.

9 examples related to the pharmacotechnical forms and the ways of application of misoprostol:

- 1) 0,05-0,20 ml gel, relatively low viscosity containing 0,9% w/v misoprostol to apply on the glans of the penis or on
- 25 the prepuce.

Synthesis:

1-1. Misoprostol	0,9 g
Hydroxypropyl Methylcellulose "3000"	2 g
Water purified to	100 ml

1-2. Misoprostol 0,9 g
Sodium Carboxymethylcellulose 2 g
Propylene Glycol 25 ml
Water purified to 100 ml

- 5 2) 0,05-0,20 ml gel of relatively high viscosity, containing 0,50% w/v in misoprostol for endourethral application at a depth 2-5 cm from the outside urethra opening.

Synthesis:

2-1. Misoprostol 0,50 g
10 Hydroxypropyl Methylcellulose "3000" 4 g
Water purified to 100 ml
2-2. Misoprostol 0,50 g
Sodium Carboxymethylcellulose 4 g
Propylene Glycol 25 ml
15 Water purified to 100 ml

- 3) 0,05-0,20 ml of aqueous solution of misoprostol containing 0,9% w/v for spreading on the glans of the penis or of the prepuce. The solution can also contain propylene glycol or glycerol in the corresponding proportions (e.g. 10%) to
20 increase the viscosity of the solution.

4) 0,05-0,20 ml of ointment or emulsion of containing 0,9% w/w misoprostol for apply on the glans of the penis or on the prepuce, where misoprostol is found spread in the continuous (aqueous) phase.

25 Synthesis:

4-1. Misoprostol 0,9 g
Vanishing Cream to 100 g

(Although for the requirements of this example as Vanishing

Cream we used Bepanthène® Cream of Roche, we have various creams o/w which are available in commerce or are described in National Pharmacopoeies and can be used for the same purpose).

- 5 5) Endourethral sticks of suitable dimensions, weight about 500 mg, containing 0,04-0,20% w/w misoprostol to apply on the urethral mucosa.

Synthesis:

5-1. Misoprostol	0,04-0,20 g
10 Glycerol	70 g
Gelatine	20 g
Water purified to	100 g

- 6) 0,05-0,25 ml gel (or more depending of response) according to the examples (1-1) and (2-1) which contains moreover 1,6%
15 w/v α -cyclodextrine.

7) 0,05-0,25 ml gel (or more depending of response) according to the example (6) which contains moreover 10 ml ethyl alcohol 96° and 0,5 mg/ml alprostadil.

Notes: 1) The incorporation of misoprostol in bases already
20 mentioned took place in normal temperature (20-25°C) and at a temperature not exceeding 40°C.

2) No significant changes in misoprostol activity was observed as a function of pH, we observed however an important reduction or/and neutralization of misoprostol action in the
25 presence of Polysorbate "80".

3) The time of appearance of the result varies from 20-40 minutes. The timing of the appearance and the intensity of the result seems to be able been positively influenced by certain

moisturising agents (e.g. Propylene Glycol, Glycerol) as well as by certain substances which reinforce by various mechanisms the transcutaneous absorption (e.g. Urea, Acid Citric).

5 High once only doses of misoprostol (>1800 mcg on the glans of the penis and >1000 mcg in the urethra) cause certain systematic undesirable effects as shudder, feeling of hardship, excitement and diarrhea. The presence of α -cyclodextrine reduces the undesirable effects and allows the application
10 once only of higher doses (>2000 mcg) without notable effect on the timing of its action but with positive effect on the intensity result and with prolonging of its duration.

5) The doses which are mentioned in the examples are only indicative since the intensity of the result depends, apart
15 from the nature and the grade of the erection dysfunction on other factors as e.g. the degree of moisturising of the underlying tissue, the physiological situation of the skin or the mucosa etc. As has already been mentioned, misoprostol is an extremely hydrophile molecule compared with other
20 prostaglandins of E₁ series (e.g. with alprostadil which can be dissolved in alcohol but its solubility in water is only 8000 mcg/100 ml at 35°C).

This consists an important advantage:

a) Because no use of organic factors is required (e.g. ethyl
25 alcohol) which usually irritate tissues and are thus unsuitable for application on the skin and especially the mucus.

b) Because it allows the incorporation of active substances

on a very small amount of excipient, suitable for application on surfaces of limited extent, as e.g. the urethra or the glans of the penis.

- 6) Because of the described irritation of the uterine fibbers
- 5 (Physicians Desc Reference, ed. Medical Economics Data, Production Company at Montrale, 48th edition, 1994, p. 2197-2199) misoprostol must not contact the female genital apparatus.

Claims

- 1) The use of misoprostol or/and its first metabolite, misoprostol acid, for the production of a drug applied topically on the glans of the penis, the prepuce or the urethra and is destined:
- a) For the symptomatic therapy of male impotence due to vascular or other causes and
- b) As aid in the diagnosis of erection dysfunction.
- 2) The use of misoprostol and/or its metabolite (misoprostol acid) according to claim (1) either as racemic mixtures or in the form of one of the stereoisomers from which they consist: [(±)-R form & (±)-S form].
- 3) The use of misoprostol and misoprostol acid according to claims (1) and (2) in the form of various galenic preparations (solutions, ointments, endourethral sticks, systems of controlled transdermal absorption) which, according to the general principles of pharmacotechnics, facilitate the application and precise administration of the right doses for the achievement of the desired therapeutical or diagnostic aim as described in claim (1).
- 4) The use of misoprostol and misoprostol acid according to claim (1), (2) and (3) in combination with other vasodilatory drugs as alprostadil for the appearance of synergistic action, as well as "passage accelerators" used normally in pharmaceutical technology aiming to increase absorption of drugs through the skin or the mucosa.
- 5) Use of misoprostol and misoprostol acid according to claim (1), (2) and (3) in combination with α-cyclodextrin or other

substances which, according to the general methods used in pharmacology impedes or retards the appearance or undesired effects of the drug or prolong their pharmaceutical action.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GR 98/00012

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K31/557

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	ANON.: "Alprostadil for erectile impotence." DRUG AND THERAP. BULL., Vol. 33, no. 8, 1995, pages 61-62, XP002049460	
A	US 5 510 384 A (MCKEE ET AL.) 23 April 1996	
A	WO 93 00894 A (SCOTT) 21 January 1993	



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

- "A" document defining the general state of the art which is not considered to be of particular relevance
- "E" earlier document but published on or after the international filing date
- "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- "O" document referring to an oral disclosure, use, exhibition or other means
- "P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

12 June 1998

Date of mailing of the international search report

22/06/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Klaver, T

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GR 98/00012

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
US 5510384	A	23-04-1996	US 5324746 A	28-06-1994
			AU 6154994 A	29-08-1994
			CA 2155971 A	18-08-1994
			EP 0682521 A	22-11-1995
			WO 9417807 A	18-08-1994
<hr/>				
WO 9300894	A	21-01-1993	AU 2300092 A	11-02-1993
			US 5708031 A	13-01-1998
<hr/>				